

AS & A Level Mathematics (9709) Paper 3

[Pure Mathematics 3]

Exam Series: February/March 2018 – May/June 2025

Format Type A:
Answers to all questions are provided as an appendix



Introduction

Each Topical Past Paper Questions Compilation contains a comprehensive collection of hundreds of questions and corresponding answer schemes, presented in worksheet format. The questions are carefully arranged according to their respective chapters and topics, which align with the latest IGCSE or AS/A Level subject content. Here are the key features of these resources:

1. The workbook covers a wide range of topics, which are organized according to the latest syllabus content for Cambridge IGCSE or AS/A Level exams.
2. Each topic includes numerous questions, allowing students to practice and reinforce their understanding of key concepts and skills.
3. The questions are accompanied by detailed answer schemes, which provide clear explanations and guidance for students to improve their performance.
4. The workbook's format is user-friendly, with worksheets that are easy to read and navigate.
5. This workbook is an ideal resource for students who want to familiarize themselves with the types of questions that may appear in their exams and to develop their problem-solving and analytical skills.

Overall, Topical Past Paper Questions Workbooks are a valuable tool for students preparing for IGCSE or AS/A level exams, providing them with the opportunity to practice and refine their knowledge and skills in a structured and comprehensive manner. To provide a clearer description of this book's specifications, here are some key details:

- Title: Cambridge AS & A Level Mathematics (9709) Paper 3 Topical Past Papers
- Subtitle: Exam Practice Worksheets With Answer Scheme
- Examination board: Cambridge Assessment International Education (CAIE)
- Subject code: 9709
- Years covered: February/March 2018 – May/June 2025
- Paper: 3
- Number of pages: 1197
- Number of questions: 532

Contents

| | | |
|----------|--|------------|
| 1 | Algebra | 7 |
| 2 | Logarithmic and exponential functions | 91 |
| 3 | Trigonometry | 143 |
| 4 | Differentiation | 221 |
| 5 | Integration | 311 |
| 6 | Numerical solution of equations | 457 |
| 7 | Vectors | 551 |
| 8 | Differential equations | 657 |
| 9 | Complex numbers | 751 |
| A | Answers | 857 |

Chapter 1

Algebra

1. 9709_m25_qp_32_Q: 9

The polynomial $6x^3 + ax^2 + bx + 9$ is denoted by $p(x)$, where a and b are constants. It is given that $(x - 3)$ is a factor of $p(x)$, and when the first derivative $p'(x)$ is divided by $(x - 3)$ the remainder is 72.

(a) Find the values of a and b . [5]

(b) When a and b have the values found in part (a), factorise $p(x)$ completely. [3]

(c) Hence solve the inequality $p(x) < 0$. [2]

2. 9709 - s25 - qp - 31 Q: 1

(a) Sketch the graph of $y = |2x - 3|$. [1]

(b) Solve the inequality $3x - 1 < |2x - 3|$. [2]

3. 9709 s25 qp 31 Q: 5

The polynomial $3x^3 + pax^2 + 7a^2x + qa^3$ is denoted by $f(x)$, where p, q and a are constants and $a \neq 0$.

When $f(x)$ is divided by $(x+2a)$ the remainder is $-22a^3$. When $f(x)$ is divided by $(3x-a)$ the remainder is $-a^3$.

Find the values of p and q .

[5]

4. 9709_s25_qp_32 Q: 2

(a) Expand $(6-x)(1-2x)^{-\frac{3}{2}}$ in ascending powers of x , up to and including the term in x^2 , simplifying the coefficients. [4]

(b) State the set of values of x for which the expansion is valid. [1]

5. 9709 s 25 qp 33 Q: 1

(a) Sketch the graph of $y = |3x - 2a|$, where a is a positive constant. [1]

(b) Hence or otherwise solve the inequality $|3x - 2a| < x + 5a$. [3]

6. 9709 s 25 qp 35 Q: 9

(a) Express $\frac{12x^2 + 55x - 2}{(3x-2)(x+6)}$ in partial fractions. [5]

(b) Hence obtain the expansion of $\frac{12x^2 + 55x - 2}{(3x-2)(x+6)}$ in ascending powers of x , up to and including the term in x^2 . [4]

7. 9709_m24_qp_32 Q: 1

Find the quotient and remainder when $x^4 - 3x^3 + 9x^2 - 12x + 27$ is divided by $x^2 + 5$. [3]

8. 9709_s24_qp_32 Q: 1

(a) Sketch the graph of $y = |x - 2a|$, where a is a positive constant. [1]

(b) Solve the inequality $2x - 3a < |x - 2a|$. [2]

9. 9709_s24_qp_32 Q: 2

Express $\frac{6x^2 - 9x - 16}{2x^2 - 5x - 12}$ in partial fractions.

[5]

10. 9709_s24_qp_33 Q: 5

Express $\frac{6x^2 - 2x + 2}{(x-1)(2x+1)}$ in partial fractions. [5]

11. 9709_w24_qp_31 Q: 1

The polynomial $4x^3 + ax^2 + 5x + b$, where a and b are constants, is denoted by $p(x)$. It is given that $(2x+1)$ is a factor of $p(x)$. When $p(x)$ is divided by $(x-4)$ the remainder is equal to 3 times the remainder when $p(x)$ is divided by $(x-2)$.

Find the values of a and b .

[5]

12. 9709_w24_qp_31 Q: 7

$$\text{Let } f(x) = \frac{5x^2 + 8x + 5}{(1 + 2x)(2 + x^2)}.$$

(a) Express $f(x)$ in partial fractions. [5]

(b) Hence find the coefficient of x^3 in the expansion of $f(x)$. [4]

13. 9709_w24_qp_33 Q: 8

Let $f(x) = \frac{7a^2}{(a-2x)(3a+x)}$, where a is a positive constant.

(a) Express $f(x)$ in partial fractions. [3]

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [4]

(c) State the set of values of x for which the expansion in part (b) is valid. [1]

14. 9709_w24_qp_33 Q: 9

(a) Find the quotient and remainder when $x^4 + 16$ is divided by $x^2 + 4$. [3]

(b) Hence show that $\int_2^{2\sqrt{3}} \frac{x^4 + 16}{x^2 + 4} dx = \frac{4}{3}(\pi + 4)$. [5]

15. 9709 - m23 - qp - 32 Q: 3

The polynomial $2x^4 + ax^3 + bx - 1$, where a and b are constants, is denoted by $p(x)$. When $p(x)$ is divided by $x^2 - x + 1$ the remainder is $3x + 2$.

Find the values of a and b .

[5]

16. 9709_s23_qp_31 Q: 2

(a) Sketch the graph of $y = |2x + 3|$. [1]

(b) Solve the inequality $3x + 8 > |2x + 3|$. [3]

17. 9709_s23_qp_32 Q: 1

Solve the inequality $|5x - 3| < 2|3x - 7|$.

[4]

18. 9709_s23_qp_33 Q: 10

$$\text{Let } f(x) = \frac{21 - 8x - 2x^2}{(1 + 2x)(3 - x)^2}.$$

(a) Express $f(x)$ in partial fractions. [5]

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

19. 9709_w23_qp_31 Q: 10

$$\text{Let } f(x) = \frac{24x + 13}{(1 - 2x)(2 + x)^2}.$$

(a) Express $f(x)$ in partial fractions. [5]

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

(c) State the set of values of x for which the expansion in (b) is valid. [1]

.....
.....
.....
.....

20. 9709_w23_qp_32 Q: 1

(a) Sketch the graph of $y = |4x - 2|$. [1]

(b) Solve the inequality $1 + 3x < |4x - 2|$.

[4]

21. 9709_w23_qp_33 Q: 3

The polynomial $2x^3 + ax^2 + bx + 6$, where a and b are constants, is denoted by $p(x)$. When $p(x)$ is divided by $(x + 2)$ the remainder is -38 and when $p(x)$ is divided by $(2x - 1)$ the remainder is $\frac{19}{2}$.

Find the values of a and b .

[5]

22. 9709_w23_qp_33 Q: 9

$$\text{Let } f(x) = \frac{17x^2 - 7x + 16}{(2 + 3x^2)(2 - x)}.$$

(a) Express $f(x)$ in partial fractions. [5]

(b) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^3 . [5]

(c) State the set of values of x for which the expansion in (b) is valid. Give your answer in an exact form. [1]

.....
.....
.....
.....
.....

23. 9709_m22_qp_32 Q: 1

Solve the inequality $|2x + 3| > 3|x + 2|$.

[4]

24. 9709 - S22 - qp - 31 Q: 2

(a) Expand $(2 - x^2)^{-2}$ in ascending powers of x , up to and including the term in x^4 , simplifying the coefficients. [4]

(b) State the set of values of x for which the expansion is valid. [1]

.....
.....
.....
.....
.....

25. 9709 S22 qp 31 Q: 5

The polynomial $ax^3 - 10x^2 + bx + 8$, where a and b are constants, is denoted by $p(x)$. It is given that $(x - 2)$ is a factor of both $p(x)$ and $p'(x)$.

(a) Find the values of a and b . [5]

(b) When a and b have these values, factorise $p(x)$ completely. [3]

Appendix A

Answers

1. 9709_m25_ms_32 Q: 9

| Question | Answer | Marks | Guidance |
|----------|--|-------|----------|
| (a) | Substitute $x = 3$ or -3 into $p(x)$ and equate to 0 or into $p'(x)$ and equate to 72 | M1* | |
| | Obtain $162 + 9a + 3b + 9 = 0$ | A1 | OE |
| | Obtain $162 + 6a + b = 72$ | A1 | OE |
| | Solve simultaneous equations to obtain either a or b after using $p(\pm 3) = 0$ and $p'(\pm 3) = 72$ | DM1 | |
| | Obtain $a = -11$ and $b = -24$ | A1 | |
| | | 5 | |

| Question | Answer | Marks | Guidance |
|----------|---|-------|---|
| (b) | Equate $(x-3)(6x^2 + Ax + B)$ to $6x^3 - 11x^2 - 24x + 9$ and obtain equations to solve for A and B | M1 | Using <i>their a</i> and <i>their b</i> . $A - 18 = a = -11, B - 3A = -b = -24,$ $-3B = 9,$ $A = 7$ and $B = -3.$ |
| | or divide $6x^3 - 11x^2 - 24x + 9$ by $x - 3$ and reach $6x^2 \pm 7x$ | | Or reach $6x^2 \pm (\text{their } a + 18)x.$ |
| | $(x-3)(6x^2 + 7x - 3)$ | A1 | SOI |
| | Obtain $(x-3)(2x+3)(3x-1)$ | A1 | |
| | | | Special Case: If only $(x-3)(x+\frac{3}{2})(x-\frac{1}{3})$ or $(x-3)(2x+3)(3x-1)$ seen, SC B1 only (but can gain two marks in (c)). |
| | | 3 | |
| (c) | Obtain one correct region $x < -\frac{3}{2}$ or $\frac{1}{3} < x < 3$ | B1 FT | Must be final answer not in working. FT is on the last two brackets (not $(x-3)$). |
| | Obtain both regions $x < -\frac{3}{2}, \frac{1}{3} < x < 3$ | B1 FT | Allow $x < -\frac{3}{2}$ and $\frac{1}{3} < x < 3.$ SC B1 for $x \leq -\frac{3}{2}, \frac{1}{3} \leq x \leq 3.$ FT is on the last two brackets (not $(x-3)$). If incorrect factor or factors in (b) but correct regions here, allow SC B1 only. |
| | | 2 | |

compiled by examinent.com

2. 9709_s25_ms_31 Q: 1

| Question | Answer | Marks | Guidance |
|----------|--|-----------|--|
| (a) | | B1 | Symmetrical. In correct position. Condone if no complete scale shown, but must see 3 and $\frac{1}{2}$ marked. Needs to exist for negative x. Must be intending straight lines. Ignore $y = 3x - 1$ if seen. |
| | | 1 | |
| (b) | Obtain critical value $\frac{4}{5}$ from $3x - 1 = 3 - 2x$ | B1 | |
| | State final answer $x < \frac{4}{5}$ | B1 | |
| | Alternative Method for Question 1(b) | | |
| | Obtain critical value $\frac{4}{5}$ from $(3x - 1)^2 = (3 - 2x)^2$ | B1 | Ignore $x = -2$ if seen. |
| | State final answer $x < \frac{4}{5}$ | B1 | |
| | | 2 | |

_____ compiled by examinent.com _____

3. 9709_s25_ms_31 Q: 5

| Question | Answer | Marks | Guidance |
|----------|--|-----------|--|
| | Use $f(-2a) = -22a^3$ | M1 | Or use long division and equate a constant remainder to $-22a^3$. |
| | Obtain $-24a^3 + 4pa^3 - 14a^3 + qa^3 = -22a^3$ | A1 | Must evaluate the terms. OE, e.g. $4p + q = 16$. |
| | Use $f\left(\frac{a}{3}\right) = -a^3$ | M1 | Or use long division and equate a constant remainder to $-a^3$. |
| | Obtain $\frac{1}{9}a^3 + \frac{1}{9}pa^3 + \frac{7}{3}a^3 + qa^3 = -a^3$ | A1 | Must evaluate the terms. OE, e.g. $p + 9q = -31$ |
| | Obtain $p = 5, q = -4$ | A1 | |
| | | 5 | |

_____ compiled by examinent.com _____

4. 9709_s25_ms_32 Q: 2

| Question | Answer | Marks | Guidance |
|----------|---|-----------|--|
| (a) | Find the first two terms of the expansion of $(1-2x)^{-\frac{3}{2}}$ | B1 | |
| | Obtain correct third term $\frac{-\frac{3}{2}(-\frac{3}{2}-1)}{2!}(-2x)^2$ or $\frac{-\frac{3}{2}(-\frac{3}{2}-1)}{2!}(2x)^2$ | B1 | $\frac{15}{2}x^2$ Ignore extra terms. |
| | Multiply <i>their</i> 3 term expansion $a+bx+cx^2$ by $(6-x)$ obtaining all necessary terms | M1 | $6+18x+45x^2-x-3x^2\dots$ Ignore extra terms. |
| | $6+17x+42x^2$ | A1 | Ignore extra terms. Allow with the terms in any order. |
| | | 4 | |
| (b) | $ x < \frac{1}{2}$ or $-\frac{1}{2} < x < \frac{1}{2}$ or $(-0.5, 0.5)$ or $]-0.5, 0.5[$ | B1 | OE B0 for an ambiguous statement. Must be strict inequality. |
| | | 1 | |

compiled by examinent.com

5. 9709_s25_ms_33 Q: 1

| Question | Answer | Marks | Guidance |
|----------|--------|-----------|--|
| (a) | | B1 | Symmetrical. In correct position. Lines intended to be straight. Must be in both first and second quadrants. Key coordinates must be correct. Ignore $y = x + 5a$ if seen. |
| | | 1 | |

| Question | Answer | Marks | Guidance |
|---|---|-------|--|
| (b) | Obtain critical value $\frac{7a}{2}$ from $x+5a=3x-2a$ | B1 | Allow if seen in an inequality. |
| | Obtain critical value $-\frac{3a}{4}$ from $x+5a=2a-3x$ | B1 | Allow if seen in an inequality. |
| | State final answer $-\frac{3a}{4} < x < \frac{7a}{2}$ | B1 | SC B1 only for $-\frac{3a}{4} < x < \frac{7a}{2}$ with <i>their a</i> from part (a). Allow any equivalent notation. Allow $-\frac{3a}{4} < x$ and $x < \frac{7a}{2}$. |
| Alternative Method for Question 1(b) | | | |
| | Solve quadratic equation $(3x-2a)^2 = (x+5a)^2$ | M1 | $8x^2 - 22ax - 21a^2 = 0$ |
| | Obtain critical values $-\frac{3a}{4}$ and $\frac{7a}{2}$ | A1 | |
| | State final answer $-\frac{3a}{4} < x < \frac{7a}{2}$ | A1 | SC B1 only for $-\frac{3a}{4} < x < \frac{7a}{2}$ with <i>their a</i> from part (a). Allow any equivalent notation. Allow $-\frac{3a}{4} < x$ and $x < \frac{7a}{2}$. |
| | | 3 | |

compiled by examinent.com

6. 9709_s25_ms_35 Q: 9

| Question | Answer | Marks | Guidance |
|---|---|-------|---|
| (a) | State or imply the form $A + \frac{B}{3x-2} + \frac{C}{x+6}$ Values for A, B and C do not need to be substituted into this form to gain full marks | B1 | $\frac{Dx+E}{3x-2} + \frac{F}{x+6}$ and $\frac{P}{3x-2} + \frac{Qx+R}{x+6}$ B0 However, can recover all marks from these $\frac{S}{3x-2} + \frac{T}{x+6}$ B0 and can only gain maximum M1A1 |
| | Use a correct method for finding a constant | M1 | |
| | Obtain one of $A = 4$, $B = 6$ and $C = -5$ or $F = -5$ or $P = 6$ or $S = 6$ or $T = -5$ | A1 | Allow maximum M1A1 for one or more 'correct' values after B0, even if from $\frac{S}{3x-2} + \frac{T}{x+6}$. $D = 12$ $E = -2$ $F = -5$ $P = 6$ $Q = 4$ $R = 19$ $S = 6$ $T = -5$ |
| | Obtain a second value from $\frac{Dx+E}{3x-2} = A + \frac{B}{3x-2}$ or $\frac{Qx+R}{x+6} = A + \frac{C}{x+6}$ | A1 | |
| | Obtain a third value | A1 | |
| Alternative Method for Question 9(a) | | | |
| | Divide numerator by denominator and reach quotient of 4 and remainder of $Px+Q$ | M1 | Or by inspection P or $Q \neq 0$ |
| | Obtain $4 + \frac{-9x+46}{(3x-2)(x+6)}$ or $4 + \frac{-9x+46}{3x^2+16x-12}$ | A1 | |
| | State or imply <i>their</i> remainder is of form $\frac{D}{3x-2} + \frac{E}{x+6}$ Values for D and E do not need to be substituted into this form to gain full marks | B1 | |

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| (a) | Obtain one of $D = 6, E = -5$ | A1 | |
| | Obtain a second value | A1 | |
| | | 5 | |
| (b) | Use the correct method to find the first two unsimplified terms of the expansion of $(3x-2)^{-1}$ or $(x+6)^{-1}$ or $(1-\frac{3}{2}x)^{-1}$ or $(1+\frac{1}{6}x)^{-1}$ | M1 | E.g. $-2^{-1} - 2^{-2}(3x)$, or $6^{-1} - 6^{-2}(x)$, or $1 + \frac{3}{2}x$ or $1 - \frac{1}{6}x$. |
| | Obtain correct unsimplified expansions up to the term in x^2 of each partial fraction $B = 6, C = -5, A = 4$ | A1FT | The FT is on B and C, e.g. $\frac{B}{-2} \left(1 + \frac{3}{2}x + \left(\frac{3}{2}x \right)^2 \right) + \frac{C}{6} \left(1 - \frac{1}{6}x + \left(\frac{1}{6}x \right)^2 \right)$ OE. |
| | $\frac{1}{6} - \frac{157}{36}x - \frac{1463}{216}x^2$ | A1 | OE Do not ISW, e.g. multiplication through by 216. Allow terms in any order. |
| | | 4 | |


compiled by examinent.com

7. 9709_m24_ms_32 Q: 1

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| | Commence division and reach partial quotient of the form $x^2 \pm 3x$ or $x^4 - 3x^3 + 9x^2 - 12x + 27 = (x^2 + 5)(Ax^2 + Bx + C) + Dx + E$ or $Ax^4 + Bx^3 + (5A + C)x^2 + 5Bx + 5C$ and reach $A = 1$ and $B = \pm 3$ | M1 | |
| | Obtain quotient $x^2 - 3x + 4$ | A1 | $A = 1, B = -3$ [$5A + C = 9$ so $C = 4$; $5B + D = -12$ so $D = 3$; $5C + E = 27$ so $E = 7$]. A pair of incorrect statements 'remainder $x^2 - 3x + 4$ ' and 'quotient $3x + 7$ ' score M1 A1 A0. |
| | Obtain remainder $3x + 7$ | A1 | |
| | $\begin{array}{r} x^2 & - & 3x & + & 4 \\ x^2 + 5 & x^4 & - & 3x^3 & + & 9x^2 - 12x + 27 \\ & & & + 5x^2 & & \\ & & - 3x^3 & + 4x^2 & & \\ & & - 3x^3 & & - 15x & \\ & & & + 4x^2 + 3x & & \\ & & & + 4x^2 & + 20 & \\ & & & + 3x & & + 7 \end{array}$ | 3 | |

compiled by examinent.com

8. 9709_s24_ms_32 Q: 1

| Question | Answer | Marks | Guidance |
|---|---|-------------|---|
| (a) |  | B1 | <p>Correct shape, roughly symmetrical. Both sections should be solid straight lines. If not drawn with a ruler the intention must be clear. Allow construction lines if dashed or clearly fainter.</p> <p>$2a$ marked on each axis (must be $2a$, not just 2).</p> <p>Needs to extend into negative x.</p> <p>If a is given a value, then B0.</p> <p>Ignore $y = 2x - 3a$ if seen.</p> |
| | | | |
| (b) | Solve linear equation or inequality to obtain critical value $x = \frac{5}{3}a$ or exact equivalent | B1 | Ignore $x = a$ if seen. |
| | Obtain $x < \frac{5}{3}a$ or exact equivalent | B1 | <p>Accept $x < \frac{10}{6}a$ or $(-\infty, \frac{5}{3}a)$. Must be strict inequality.</p> <p>Need a clear final solution: $x > a$ or $x < a$ must be rejected if seen as part of the working. Rejection can be implied, e.g. if only the correct inequality is underlined.</p> <p>B0 B0 if a is given a value.</p> |
| Alternative Method for Question 1(b) | | | |
| Solve quadratic equation $(2x - 3a)^2 = (x - 2a)^2$ to obtain critical value $x = \frac{5}{3}a$ or exact equivalent | | (B1) | $(3x^2 - 8ax + 5a^2 = 0)$ <p>Ignore $x = a$ if seen.</p> |
| Obtain $x < \frac{5}{3}a$ or exact equivalent | | (B1) | <p>Accept $x < \frac{10}{6}a$ or $(-\infty, \frac{5}{3}a)$. Must be strict inequality.</p> <p>Need a clear final solution: $x > a$ or $x < a$ must be rejected if seen as part of the working. Rejection can be implied, e.g. if only the correct inequality is underlined.</p> <p>B0 B0 if a is given a value.</p> |
| | | 2 | |

compiled by examinent.com

9. 9709_s24_ms_32 Q: 2

| Question | Answer | Marks | Guidance |
|----------|--|-----------|---|
| | State or imply the form $A + \frac{B}{2x+3} + \frac{C}{x-4}$ | B1 | $\frac{Dx+E}{2x+3} + \frac{F}{x-4}$ and $\frac{P}{2x+3} + \frac{Qx+R}{x-4}$ are also valid. |
| | Use a correct method for finding a constant | M1 | SC: If score B0, they can score M1 A1 for one correct constant. B0 M1 A0 available if they substitute two values to form simultaneous equations but get an incorrect answer, or they substitute one value and make an arithmetic error. |
| | Obtain one of $A = 3$, $B = -2$ and $C = 4$ | A1 | SC: If the horizontal equation is correct apart from an incorrect value for A , the other A marks may be available. |
| | Obtain a second value | A1 | SC: If denominator factorised as $(x + \frac{3}{2})(x - 4)$ can score a maximum of B0 M1 A1 A1 A0 for a split involving 3 terms. |
| | Obtain a third value | A1 | ISW Statement of the final split is not required. |

| Question | Answer | Marks | Guidance |
|--|--------|---|----------|
| Alternative method for Question 2 | | | |
| Divide numerator by denominator | (M1) | | |
| Obtain $3 + \frac{Px+Q}{2x^2-5x-12}$ | (A1) | $\left(3 + \frac{6x+20}{(2x+3)(x-4)} \right)$ | |
| State or imply the form $\frac{Px+Q}{2x^2-5x-12} = \frac{D}{2x+3} + \frac{E}{x-4}$ | (B1) | Must deal with the 3 separately or include it correctly on both sides in their split. | |
| Obtain one of $D = -2$ and $E = 4$ | (A1) | SC: If denominator factorised as $(x + \frac{3}{2})(x - 4)$, then can score a maximum of B0 M1 A1 A1 A0 for a split involving three terms. | |
| Obtain a second value | (A1) | ISW Statement of the final split is not required. | |
| | | 5 | |

compiled by examinent.com

10. 9709_s24_ms_33 Q: 5

| Question | Answer | Marks | Guidance |
|--|--------|---|----------|
| Alternative Method for Question 5 | | | |
| State or imply the form $A + \frac{B}{x-1} + \frac{C}{2x+1}$ | (B1) | | |
| Use a correct method for finding a constant | (M1) | Correct appropriate method. | |
| Obtain one of $A = 3$, $B = 2$ and $C = -3$ | (A1) | | |
| Obtain a second value | (A1) | | |
| Obtain a third value | (A1) | | |
| Alternative Method for Question 5 | | | |
| Divide numerator by denominator to reach $A = 3$ | (M1) | May be implied by 3 [+] $\frac{ax+b}{(x-1)(2x+1)}$ with a and b not both 0. | |
| Obtain $3 + \frac{x+5}{(x-1)(2x+1)}$ | (A1) | | |
| State or imply the form $\frac{D}{x-1} + \frac{E}{2x+1}$ | (B1) | | |
| Obtain one of $D = 2$ and $E = -3$ | (A1) | | |
| Obtain a second value | (A1) | | |
| | | 5 | |

compiled by examinent.com

11. 9709_w24_ms_31 Q: 1

| Question | Answer | Marks | Guidance |
|----------|--|-----------|---|
| | Substitute $x = -\frac{1}{2}$ and equate the result to zero | M1 | |
| | Obtain a correct equation, e.g. $-\frac{4}{8} + \frac{a}{4} - \frac{5}{2} + b = 0$ | A1 | $(\frac{a}{4} + b = 3)$ Any equivalent form. |
| | Substitute $x = 2$ and $x = 4$ and use $p(4) = 3p(2)$ | M1 | If using long division, M1 is for correct use of two constant remainders. Condone if 3 is on the wrong side. |
| | Obtain a correct equation, e.g. $3(32 + 4a + 10 + b) = 256 + 16a + 20 + b$ | A1 | $(-2a + b = 75)$ Any equivalent form. |
| | Obtain $a = -32$ and $b = 11$ | A1 | |
| | | 5 | |

compiled by examinent.com

12. 9709_w24_ms_31 Q: 7

| Question | Answer | Marks | Guidance |
|----------|---|--------------|---|
| (a) | State or imply the form $\frac{A}{1+2x} + \frac{Bx+C}{2+x^2}$ | B1 | |
| | Use a correct method to find a constant | M1 | |
| | Obtain one of $A = 1$, $B = 2$ and $C = 3$ | A1 | |
| | Obtain a second value | A1 | |
| | Obtain the third value | A1 | |
| | | 5 | |
| (b) | State $\frac{-1 \cdot -2 \cdot -3}{3!} (2x)^3$ or -8 | B1 FT | Correct term in x^3 or coefficient of x^3 in the expansion of $A(1+2x)^{-1}$. Any equivalent form. |
| | Use a correct method to obtain the coefficient of x^2 in the expansion of $(2+x^2)^{-1}$ or the coefficient of x^2 in the expansion of $(1+\frac{x^2}{2})^{-1}$. | M1 | Do not need to deal with 2^{-1} at this stage. |
| | Obtain $(Bx+C) \times \frac{1}{2} \times -\frac{1}{2}x^2$ or $-\frac{B}{4}x^3$ or $-\frac{B}{4}$ | A1 FT | Follow <i>their B</i> (and <i>C</i>). |
| | Obtain final answer $-8\frac{1}{2}$ or $-8\frac{1}{2}x^3$ | A1 | Or simplified equivalent. Ignore additional terms for other powers of x . |
| | | 4 | |

compiled by examinent.com

13. 9709_w24_ms_33 Q: 8

| Question | Answer | Marks | Guidance |
|----------|---|-------|---|
| (a) | State or imply the form $\frac{A}{a-2x} + \frac{B}{3a+x}$ and use a correct method to find a constant | M1 | |
| | Obtain $A=2a$ or $B=a$ | A1 | |
| | Obtain $A=2a$ and $B=a$ | A1 | |
| | | 3 | |
| (b) | Use a correct method to obtain the first two terms of the expansion of $(a-2x)^{-1}$ or $\left(1-\frac{2x}{a}\right)^{-1}$ or $(3a+x)^{-1}$ or $\left(1+\frac{x}{3a}\right)^{-1}$ | M1 | |
| | Obtain $+2\left(1+\frac{2x}{a} + \frac{4x^2}{a^2} + \dots\right)$ | A1ft | OE. May be unsimplified. Follow <i>their A, B</i> for an expansion involving a . |
| | Obtain $+\frac{1}{3}\left(1-\frac{x}{3a} + \frac{x^2}{9a^2} + \dots\right)$ | A1ft | OE. May be unsimplified. Follow <i>their A, B</i> for an expansion involving a . |
| | Obtain $+\frac{7}{3} + \frac{35x}{9a} + \frac{217x^2}{27a^2}$ | A1 | Or simplified equivalent. Final answer. Ignore any terms in higher powers of x . Do not ISW, e.g. multiplying by $27a^2$. Condone different order of terms. |

| Question | Answer | Marks | Guidance |
|----------|---|-------|---|
| (b) | Alternative Method for Question 8(b) | | |
| | Expanding $7a^2(a-2x)^{-1}(3a+x)^{-1}$ from the original question. | M1 | |
| | Use a correct method to obtain the first two terms of the expansion of $(a-2x)^{-1}$ or $\left(1-\frac{2x}{a}\right)^{-1}$ or $(3a+x)^{-1}$ or $\left(1+\frac{x}{3a}\right)^{-1}$ | | |
| | Obtain $+7a\left(1+\frac{2x}{a} + \frac{4x^2}{a^2} + \dots\right)$ or $+\frac{7a}{3}\left(1-\frac{x}{3a} + \frac{x^2}{9a^2} + \dots\right)$ | A1 | OE. May be unsimplified. May be implied by the expression shown for the next A1. |
| | Obtain $+\frac{7}{3}\left(1+\frac{2x}{a} + \frac{4x^2}{a^2} + \dots\right)\left(1-\frac{x}{3a} + \frac{x^2}{9a^2} + \dots\right)$ | A1 | OE. May be unsimplified. |
| | Obtain $+\frac{7}{3} + \frac{35x}{9a} + \frac{217x^2}{27a^2}$ | A1 | Or simplified equivalent. Final answer. Ignore any terms in higher powers of x . Do not ISW, e.g. multiplying by $27a^2$. Condone different order of terms. |
| (c) | | 4 | |
| | $ x < \frac{a}{2}$ | B1 | Or $-\frac{a}{2} < x < \frac{a}{2}$. Mark final answer. Must make a clear statement. |
| | | 1 | |

14. 9709_w24_ms_33 Q: 9

| Question | Answer | Marks | Guidance |
|---|-------------------------------------|-------|---|
| (a) | Divide to obtain quotient $x^2 + k$ | M1 | k is a constant. |
| | Obtain quotient $x^2 - 4$ | A1 | If quotient stated separately, mark at this stage. |
| | Obtain remainder 32 | A1 | If remainder stated separately, mark at this stage. Need not state which is quotient and remainder, but if stated wrongly, max 2/3. After a correct division, still allow the marks if then written as $x^2 - 4 + \frac{32}{x^2 + 4}$. |
| Alternative Method for Question 9(a) | | | |
| | Expands brackets to get $B = 0$ | M1 | $(x^2 + 4)(x^2 + Bx + C) + D = x^4 + Bx^3 + (C + 4)x^2 + 4Bx + 4C + D$ |
| | $C = -4$ | A1 | |
| | $D = 32$ | A1 | Need not state which is quotient and remainder, but if stated wrongly, max 2/3. |
| | | 3 | |

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| (b) | $\frac{1}{3}x^3 - 4x$ | B1 FT | Follow <i>their</i> quotient of form $Ax^2 + B$. |
| | Obtain $p \tan^{-1} qx$ where $q = 2$ or $q = \frac{1}{2}$ | M1 | |
| | Obtain $16 \tan^{-1} \frac{1}{2}x$ | A1 FT | Follow <i>their</i> constant remainder, i.e. $\left(\frac{\text{their constant remainder}}{2}\right) \tan^{-1} \frac{1}{2}x$. |
| | Use limits correctly in an expression containing $p \tan^{-1} qx$ where $q = 2$ or $q = \frac{1}{2}$ and $rx^3 + sx$ | M1 | Terms need not be evaluated, e.g. $\left[8\sqrt{3} - 8\sqrt{3}\right] + 16 \tan^{-1} \sqrt{3} - \left(\frac{8}{3} - 8 + 16 \tan^{-1} 1\right)$ or $\frac{8}{3} - 8$ can be $-\frac{16}{3}$, $16 \tan^{-1} \sqrt{3}$ can be $\frac{16\pi}{3}$, $16 \tan^{-1} 1$ can be 4π . |
| | Obtain $\frac{4}{3}(\pi + 4)$ from full and correct working | A1 | AG |
| | | 5 | |

compiled by examinent.com

15. 9709_m23_ms_32 Q: 3

| Question | Answer | Marks | Guidance |
|----------|--|-------|---|
| | Commence division and reach partial quotient $2x^2 + (a \pm 2)x$ | M1 | $ \begin{aligned} 2x^2 + (a+2)x + a &\quad \text{need } 2x^2 + (a \pm 2)x \\ (x^2 - x + 1) 2x^4 + ax^3 + 0x^2 + bx - 1 & \\ 2x^4 - 2x^3 + 2x^2 & \\ (a+2)x^3 - 2x^2 + bx & \\ (a+2)x^3 - (a+2)x^2 + (a+2)x & \\ ax^2 + (b - (a+2))x - 1 & \\ ax^2 - ax + a & \\ (b-2)x - (1+a) & \\ 3x + 2 & \end{aligned} $ <p>Working backwards from remainder: $2x^2 + (\dots)x + 3 \text{ M1 } 2x^2 - x - 3 \text{ A1}$</p> |
| | Obtain correct quotient $2x^2 + (a+2)x + a$ | A1 | Allow sign error e.g. in $b - 2$. |
| | Set <i>their</i> linear remainder equal to part of “ $3x + 2$ ” and solve for a or for b | M1 | Remainder = $3x + 2 = (b-2)x - 1 - a$. Allow for just equating x term or constant term. |
| | Obtain answer $a = -3$ | A1 | |
| | Obtain answer $b = 5$ | A1 | |

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| | Alternative method for Question 3 | | |
| | State $2x^4 + ax^3 + 0x^2 + bx - 1 = (x^2 - x + 1)(2x^2 + Ax + B) + 3x + 2$ and form and solve equation(s) to obtain A or B | M1 | e.g. $0 = B - A + 2$ and $-1 = B + 2$. |
| | Obtain $A = -1$, $B = -3$ | A1 | |
| | Form and solve equations for a or for b | M1 | e.g. $a = A - 2$ or $b = -B + A + 3$. |
| | Obtain answer $a = -3$ | A1 | |
| | Obtain answer $b = 5$ | A1 | |
| | Alternative method for Question 3 | | |
| | Use remainder theorem with $x = \frac{1 \pm \sqrt{-3}}{2}$ or $x = \frac{1 \pm i\sqrt{3}}{2}$ | M1 | Allow for correct use of a reasonable attempt at either root in exact or decimal form in the remainder theorem $x^2 = \frac{-1 + \sqrt{-3}}{2} \quad x^3 = -1 \quad x^4 = \frac{-1 - \sqrt{-3}}{2}$. |
| | Obtain $-a + \frac{b}{2} \pm \frac{b\sqrt{-3}}{2} \mp \sqrt{-3} - 2 = \frac{7}{2} \pm \frac{3\sqrt{-3}}{2}$ or $-a + \frac{b}{2} \pm \frac{bi\sqrt{3}}{2} \mp i\sqrt{3} - 2 = \frac{7}{2} \pm \frac{3i\sqrt{3}}{2}$ | A1 | Expand brackets and obtain exact equation for either root. Accept exact equivalent. |
| | Solve simultaneous equations, or single equation, for a or for b | M1 | |
| | Obtain answer $a = -3$ from exact working | A1 | |
| | Obtain answer $b = 5$ from exact working | A1 | |
| | | 5 | |

compiled by examinent.com

16. 9709_s23_ms_31 Q: 2

| Question | Answer | Marks | Guidance |
|----------|--------|-----------|--|
| (a) | | B1 | Show a recognizable sketch graph of $y = 2x + 3 $. (Ignore any attempt to sketch $y = 3x + 8$). Straight lines. Vertex in approximately correct position on x axis. Symmetry. |

1

| Question | Answer | Marks | Guidance |
|-----------------------------|--|-------------|--|
| (b) | Find x-coordinate of intersection with $y = 3x + 8$ | M1 | |
| | Obtain $x = -\frac{11}{5}$ | A1 | |
| | State final answer $x > -\frac{11}{5}$ only | A1 | $(x > -2.2)$ Do not condone \geq for $>$. |
| Alternative Method 1 | | | |
| | Solve the linear inequality $3x + 8 > -(2x + 3)$, or corresponding linear equation | M1 | |
| | Obtain critical value $x = -\frac{11}{5}$ | A1 | |
| | State final answer $x > -\frac{11}{5}$ only | A1 | $(x > -2.2)$ Do not condone \geq for $>$. |
| Alternative Method 2 | | | |
| | Solve the quadratic inequality $(3x + 8)^2 > (2x + 3)^2$, or corresponding quadratic equation | (M1) | $5x^2 + 36x + 55$. |
| | Obtain critical value $x = -\frac{11}{5}$ | (A1) | Ignore -5 if seen. |
| | State final answer $x > -\frac{11}{5}$ only | (A1) | $(x > -2.2)$ Do not condone \geq for $>$. |
| | | 3 | |

compiled by examinent.com

17. 9709_s23_ms_32 Q: 1

| Question | Answer | Marks | Guidance |
|--|--|-----------|--|
| | State or imply non-modular inequality $(5x-3)^2 < 2^2(3x-7)^2$, or corresponding quadratic equation, or pair of linear equations $(5x-3) = \pm 2(3x-7)$ | B1 | $11x^2 - 138x + 187 > 0$. |
| | Solve a 3-term quadratic, or solve two linear equations for x | M1 | If no working is shown, the M1 is implied by the correct roots for an incorrect quadratic. |
| | Obtain critical values $x = \frac{17}{11}$ and $x = 11$ | A1 | Accept 1.55 or better. |
| | State final answer $x < \frac{17}{11}$, $x > 11$ | A1 | Strict inequality required. In set notation, allow notation for open sets but not for closed sets e.g. accept $(-\infty, \frac{17}{11}) \cup (11, \infty)$ or $(-\infty, \frac{17}{11}) \cup [11, \infty)$ but not $(-\infty, \frac{17}{11}] \cup [11, \infty)$. Allow 'or' but not 'and'. Accept \cup . Final A0 for $\frac{17}{11} > x > 11$. Exact values expected but ISW if exact inequalities seen followed by decimal approx. |
| Alternative Method for Question 1 | | | |
| | Obtain critical value $x = 11$ from a graphical method, or by inspection, or by solving a linear equation or an inequality | B1 | |
| | Obtain critical value $x = \frac{17}{11}$ similarly | B2 | Accept decimal value. |
| | State final answer $x < \frac{17}{11}$, $x > 11$ | B1 | Strict inequality required. See notes above. |
| | | 4 | |

compiled by examinent.com

18. 9709_s23_ms_33 Q: 10

| Question | Answer | Marks | Guidance |
|----------|--|-----------|---|
| (a) | State or imply the form $\frac{A}{1+2x} + \frac{B}{3-x} + \frac{C}{(3-x)^2}$ | B1 | Alternative form: $\frac{A}{1+2x} + \frac{Dx+E}{(3-x)^2}$. |
| | Use a correct method to find a constant | M1 | Incorrect format for partial fractions: Allow M1 and a possible A1 if obtain one of these correct values. Max 2/5 Allow M1 even if multiply up by $(1+2x)(3-x)^3$. |
| | Obtain one of $A = 2$, $B = 2$ and $C = -3$ | A1 | Alternative form: obtain one of $A = 2$, $D = -2$ and $E = 3$. |
| | Obtain a second value | A1 | |
| | Obtain the third value | A1 | Do not need to substitute values back into original form. |
| | | 5 | If $\frac{A}{1+2x} + \frac{B}{3-x} + \frac{Cx+D}{(3-x)^2}$ B0 but M1 A1 for A , A1 for B and A1 for C and D . If $C = 0$ then recovers B1 from above. |

| Question | Answer | Marks | Guidance |
|----------|--|-------------------------|---|
| (b) | Use a correct method to obtain the first two terms of one of the unsimplified expansions $(1+2x)^{-1}, \left(1-\frac{1}{3}x\right)^{-1}, \left(1-\frac{1}{3}x\right)^{-2}, (3-x)^{-1}, (3-x)^{-2}$ | M1 | $(1+2x)^{-1} = 1 + (-1)(2x) + \dots$ $\left(1-\frac{1}{3}x\right)^{-1} = 1 + (-1)(-x/3) + \dots$ $\left(1-\frac{1}{3}x\right)^{-2} = 1 + (-2)(-x/3) + \dots$ $(3-x)^{-1} = 3^{-1} + (-1)3^{-2}(-x) \dots$ $(3-x)^{-2} = 3^{-2} + (-2)3^{-3}(-x) + \dots$ |
| | Obtain the correct unsimplified expansions up to the term in x^2 for each partial fraction If correct, should be working with $\frac{2}{1+2x} + \frac{2}{3-x} - \frac{3}{(3-x)^2}$ or $\frac{2}{1+2x} + \frac{-2x+3}{(3-x)^2}$ | A1 FT A1 FT A1 FT | Follow through on <i>their</i> A, B, C $A(1 + (-1)(2x) + ((-1)(-2)/2)(2x)^2 + \dots)$ $\frac{B}{3} (1 + (-1)(-x/3) + ((-1)(-2)/2)(-x/3)^2 + \dots)$ $\frac{C}{3^2} (1 + (-2)(-x/3) + ((-2)(-3)/2)(-x/3)^2 + \dots)$. Must be <i>their</i> coefficients from (a) but may be unsimplified expansions for FT marks. If correct, expect to see $2(1 - 2x + (2x)^2)$ or $2 - 4x + 8x^2$ $\frac{2}{3} (1 + \frac{x}{3} + \left(\frac{x}{3}\right)^2)$ or $\frac{2}{3} + \frac{2}{9}x + \frac{2}{27}x^2$ $-\frac{1}{3} (1 + \frac{2x}{3} + (3)\left(\frac{x}{3}\right)^2)$ or $-\frac{1}{3} - \frac{2}{9}x - \frac{x^2}{9}$. |
| | Obtain final answer $\frac{7}{3} - 4x + \frac{215}{27}x^2$ | A1 | Accept $2\frac{1}{3} - 4x + 7\frac{26}{27}x^2$. No ISW. |

| Question | Answer | Marks | Guidance |
|----------|--|--------------|--|
| (b) | Alternative Method for Question 10(b) | | |
| | For the form $\frac{A}{1+2x} + \frac{Dx+E}{(3-x)^2}$ | M1* | For the first two terms of an expanded partial fraction, following <i>their</i> A, D, E. |
| | Obtain the correct unsimplified expansions up to the term in x^2 for each partial fraction | A1FT A1FT | $A(1 + (-1)(2x) + ((-1)(-2)/2)(2x)^2 + \dots) +$ $(Dx+E)\frac{1}{3^2} (1 + (-2)(-x/3) + ((-2)(-3)/2)(-x/3)^2 +)$ $2(1 - 2x + (2x)^2 + \dots)$ $+ \frac{-2x+3}{3^2} (1 + \frac{2x}{3} + (3)\left(\frac{x}{3}\right)^2 + \dots)$. |
| | Multiply out fully | DM1 | Provided $DE \neq 0$. Ignore cubic terms and above. Allow error in one term but all terms must be present. If correct, expect to see $2 - 4x + 8x^2 - \frac{2}{9}x - \frac{4}{27}x^2 + \frac{1}{3} + \frac{2}{9}x + \frac{1}{9}x^2$ |
| | Obtain final answer $\frac{7}{3} - 4x + \frac{215}{27}x^2$ | A1 | Accept $2\frac{1}{3} - 4x + 7\frac{26}{27}x^2$. No ISW |

| Question | Answer | Marks | Guidance |
|----------|---|-------|---|
| (b) | Alternative Method for Question 10(b): Maclaurin's Series | | |
| | Correct derivatives for $A(1+2x)^{-1}$, $B(3-x)^{-1}$ and $C(3-x)^{-2}$ $(-1)(2)A(1+2x)^{-2}$, $(-1)(-1)B(3-x)^{-2}$ and $(-2)(-1)C(3-x)^{-3}$ | B1 FT | |
| | One of following $(-2)(2)(-1)(2)A(1+2x)^{-3}$, $(-2)(-1)(-1)(-1)B(3-x)^{-3}$ and $(-3)(-1)(-2)(-1)C(3-x)^{-4}$ | B1 FT | |
| | All correct | B1 FT | |
| | Substitute in $f(0) + xf'(0) + \frac{x^2}{2} f''(0)$ | M1 | |
| | Obtain final answer $\frac{7}{3} - 4x + \frac{215}{27}x^2$ | A1 | Accept $2\frac{1}{3} - 4x + 7\frac{26}{27}x^2$. No ISW |
| | | 5 | |

19. 9709_w23_ms_31 Q: 10

| Question | Answer | Marks | Guidance |
|----------|--|-------|---|
| (a) | State or imply the form $\frac{A}{1-2x} + \frac{B}{2+x} + \frac{C}{(2+x)^2}$ | B1 | |
| | Use a correct method for finding a coefficient | M1 | $A(2+x)^2 + B(1-2x)(2+x) + C(1-2x) = 24x + 13.$ |
| | Obtain one of $A = 4$, $B = 2$ and $C = -7$ | A1 | If errors in equating still allow A marks for A and C . |
| | Obtain a second value | A1 | |
| | Obtain the third value | A1 | Mark the form $\frac{A}{1-2x} + \frac{Dx+E}{(2+x)^2}$, where $A = 4$, $D = 2$ and $E = -3$, B1 M1 A1 A1 A1 as above. If there are extra term in partial fractions, that is 4 unknowns A , B , D and E then B0 unless recover at end, e.g. by setting $B = 0$. If B set to any value other than 0 and all coefficients correctly found to their new values then allow all A marks, but still B0 for partial fraction expression. Hence A1 for each coefficient, but nothing for coefficient set to specific value. Another case of extra term in partial fraction expression, namely $+F$, mark as above but need $F = 0$ to recover B1. |
| | | 5 | |

| Question | Answer | Marks | Guidance |
|----------|---|-------|---|
| (b) | Use a correct method to find the first two terms of the expansion of $(1-2x)^{-1}$, $(2+x)^{-1}$, $(2+x)^{-2}$, $\left(1+\frac{x}{2}\right)^{-1}$ or $\left(1+\frac{x}{2}\right)^{-2}$ | M1 | Symbolic coefficients are not sufficient for the M1. |
| | Obtain correct un-simplified expansions up to the term in x^2 of each partial fraction | A1 FT | $A\left(1+(-1)(-2x)+\frac{(-1)(-2)}{2}(-2x)^2+\dots\right) A = 4.$ |
| | | A1 FT | $\frac{B}{2}\left(1+(-1)\left(\frac{x}{2}\right)+\frac{(-1)(-2)}{2}\left(\frac{x}{2}\right)^2+\dots\right) B = 2.$ |
| | | A1 FT | $\frac{C}{4}\left(1+(-2)\left(\frac{x}{2}\right)+\frac{(-2)(-3)}{2}\left(\frac{x}{2}\right)^2+\dots\right) C = -7$ $= 4(1+2x+4x^2) + 2/2(1-x/2+x^2/4) - 7/4(1-x+3x^2/4)$ $= (4+1-7/4) + (8-1/2+7/4)x + (16+1/4-21/16)x^2$ The FT is on A , B , C . |

| Question | Answer | Marks | Guidance |
|----------|--|-----------|--|
| | Obtain final answer $\frac{13}{4} + \frac{37}{4}x + \frac{239}{16}x^2$ | A1 | <p>OE $(Dx + E)/4 [1 + (-2)(x/2) + (-2)(-3)(x/2)^2/2 \dots]$ $D = 2 \quad E = -3$ The FT is on A, D, E.</p> <p>MacLaurin's Series $f(0) = 13/4 \quad B1 \quad f'(0) = 37/4 \quad B1 \quad f''(0) = 239/8 \quad B1$. $\frac{13}{4} + \frac{37}{4}x + \frac{239}{8}x^2/2$ or equivalent M1 A1.</p> <p>If $1 + \frac{37}{4}x + \frac{239}{8}x^2/2$ then M0 A0 unless <i>their</i> $f(0)$ actually is 1.</p> <p>For the A, D, E form of fractions, give M1 A1FT A1FT for the expanded partial fractions, then, if $D \neq 0$, M1 for multiplying out fully, and A1 for the final answer.</p> <p>If final answer has been multiplied throughout (e.g. by 16) then A0 at the end</p> |
| | | 5 | |
| 10(c) | $ x < \frac{1}{2}$ | B1 | OE |
| | | 1 | |

compiled by examinent.com

20. 9709_w23_ms_32 Q: 1

| Question | Answer | Marks | Guidance |
|----------|--------|-----------|--|
| (a) | | B1 | <p>Show a recognizable sketch graph of $y = 4x - 2$. Roughly symmetrical. Should extend into the second quadrant. Ignore $y = 4x - 2$ below the axis if intention is clear e.g. dashed or the required lines are clearly bolder. Some indication of scale on both axes – accept dashes. Must go beyond $(0, 2)$ and $(1, 2)$. Ignore any attempt to sketch $y = 1 + 3x$.</p> |
| | | 1 | |

| Question | Answer | Marks | Guidance |
|---|--|-----------|--|
| (b) | Obtain critical value $x=3$ | B1 | Allow incorrect inequality. Allow if later rejected. Allow $\frac{21}{7}$. |
| | Solve the linear equation $1+3x=2-4x$ | M1 | Or corresponding linear inequality. |
| | Obtain critical value $\frac{1}{7}$ | A1 | Allow 0.143 or better. Allow incorrect inequality. Allow if later rejected. |
| | Obtain final answer $x < \frac{1}{7}$ [or] $x > 3$ | A1 | Or equivalent. Allow with a comma, or nothing between. Strict inequalities only. Exact values. A0 for $\frac{1}{7} > x > 3$ A0 for $x < \frac{1}{7}$ and $x > 3$. |
| Alternative method for question 1(b) | | | |
| | Solve the quadratic inequality $(4x-2)^2 > (1+3x)^2$, or corresponding quadratic equation | M1 | e.g. $7x^2 - 22x + 3 = 0$. Available if they start with the correct equation / inequality, have a correct method for squaring (i.e. not $(a+b)^2 = a^2 + b^2$) and a correct method for solving. Need to obtain at least one critical value. |
| | Obtain critical value $x=3$ | A1 | Allow incorrect inequality. Allow if later rejected. Allow $\frac{21}{7}$. |
| | Obtain critical value $\frac{1}{7}$ | A1 | Allow 0.143 or better. Allow incorrect inequality. Allow if later rejected. |
| | Obtain final answer $x < \frac{1}{7}$ [or] $x > 3$ | A1 | Or equivalent. Strict inequalities only. Allow with a comma, or nothing between. Exact values. A0 for $\frac{1}{7} > x > 3$ A0 for $x < \frac{1}{7}$ and $x > 3$. |
| | | 4 | |

_____ compiled by examinent.com _____

21. 9709_w23_ms_33 Q: 3

| Question | Answer | Marks | Guidance |
|----------|--|-----------|--|
| | $2(-2)^3 + a(-2)^2 + b(-2) + 6 = -38$ Allow errors $\begin{array}{r} x+2 \underline{\quad 2x^2 + (a-4)x + b - 2a + 8} \\ \quad 2x^3 + ax^2 \quad \quad \quad bx \quad \quad \quad + 6 \\ \underline{2x^3 + 4x^2} \\ \quad (a-4)x^2 + bx \\ \quad (a-4)x^2 + (2a-8)x \\ \quad (b-2a+8)x + 6 \\ \quad (b-2a+8)x + 2b - 4a + 16 \\ \hline \quad \quad \quad \quad \quad 4a - 2b - 10 \end{array}$ | M1 | Substitute $x = -2$ and equate the result to -38 or divide by $x + 2$ to obtain quadratic quotient, and equate constant remainder to -38 . |

| Question | Answer | Marks | Guidance |
|----------|--|-------|---|
| | Obtain a correct evaluated equation, e.g. $-16 + 4a - 2b + 6 = -38$ or $4a - 2b = -28$ | A1 | |
| | $2\left(\frac{1}{2}\right)^3 + a\left(\frac{1}{2}\right)^2 + b\left(\frac{1}{2}\right) + 6 = \frac{19}{2}$ Allow errors $\begin{array}{r} x^2 + a+1 \\ 2x-1 \end{array} \overline{) 2x^3 + ax^2 + bx + 6}$ $\begin{array}{r} 2x^3 - x^2 \\ (a+1)x^2 + bx \end{array}$ $\begin{array}{r} (a+1)x^2 - \left(\frac{a}{2} + \frac{1}{2}\right)x \\ \hline \left(b + \frac{a}{2} + \frac{1}{2}\right)x + 6 \end{array}$ $\begin{array}{r} \left(b + \frac{a}{2} + \frac{1}{2}\right)x - \left(\frac{b}{2} + \frac{a}{4} + \frac{1}{4}\right) \\ \hline 6 + \frac{b}{2} + \frac{a}{4} + \frac{1}{4} \end{array}$ | M1 | Substitute $x = \frac{1}{2}$ and equate the result to $\frac{19}{2}$ or divide by $2x - 1$ to obtain quadratic quotient, and equate constant remainder to $\frac{19}{2}$. |
| | Obtain a correct evaluated equation, e.g. $\frac{1}{4} + \frac{a}{4} + \frac{b}{2} + 6 = \frac{19}{2}$ or $\frac{a}{4} + \frac{b}{2} = \frac{13}{4}$ | A1 | |
| | Obtain $a = -3$ and $b = 8$ | A1 | ISW |
| | | 5 | |

compiled by examinent.com

22. 9709_w23_ms_33 Q: 9

| Question | Answer | Marks | Guidance |
|----------|---|-------|--|
| (a) | State or imply the form $\frac{Ax+B}{2+3x^2} + \frac{C}{2-x}$ | B1 | If incorrect partial fractions e.g. $A = 0$ or $Ax^2 + B$ then M1, A1 A0 for correct C . Only allow single A1 even if other coefficients correct. B1 recoverable by a correct form end statement. |
| | Use a correct method for finding a coefficient | M1 | e.g. $(Ax+B)(2-x) + C(2+3x^2) = (3C-A)x^2 + (2A-B)x + (2B+2C) = 17x^2 - 7x + 16$ |
| | Obtain one of $A = -2$, $B = 3$ and $C = 5$ | A1 | If error present in above still allow A1 for C . |
| | Obtain a second value | A1 | |
| | Obtain the third value | A1 | Extra term in partial fractions, $D/(2+3x^2)$, that is 4 unknowns A , B , C and D then B0 unless recover at end, e.g. by setting B or $D = 0$. If B or D set to any value other than 0 and all coefficients correctly found to their new values then allow all A marks, but still B0 for partial fraction expression unless $B + D$ combined. Hence A1 for each coefficient, but nothing for coefficient set to specific value. Another case of extra term in partial fraction expression, namely $+ F$, mark as above but need $F = 0$ to recover B1. |
| | | 5 | |

| Question | Answer | Marks | Guidance |
|----------|--|-------|--|
| (b) | Use a correct method to find the first two terms of the expansion $(2-x)^{-1} = 2^{-1} + (-1) 2^{-2}(-x) + [(-1)(-2)2^{-3}(-x)^2/2!]$, $\left(1 + \frac{3x^2}{2}\right)^{-1} = 1 - \frac{3x^2}{2} \quad \text{or} \quad \left(1 - \frac{x}{2}\right)^{-1} = 1 - \left(\frac{-x}{2}\right)$ | M1 | Symbolic coefficients are not sufficient for the M1. |
| | $\frac{Ax+B}{2} \left[1 + \left(-1\right) \frac{3x^2}{2} \dots\right] \quad A = -2 \quad B = 3$ $\frac{C}{2} \left[1 + \left(-1\right) \left(\frac{-x}{2}\right) + \frac{(-1)(-2)}{2} \left(\frac{-x}{2}\right)^2 \dots\right] \quad C = 5$ | A1 FT | Obtain correct un-simplified expansions up to the term in x^3 of each partial fraction. |
| | $= \frac{3-2x}{2} \left(1 - \frac{3x^2}{2}\right) + \frac{5}{2} \left(1 + \frac{x}{2} + \frac{x^2}{4} + \frac{x^3}{8}\right)$ $= \left(\frac{3}{2} + \frac{5}{2}\right) + \left(-1 + \frac{5}{4}\right)x + \left(-\frac{9}{4} + \frac{5}{8}\right)x^2 + \left(\frac{3}{2} + \frac{5}{16}\right)x^3$ | A1 FT | Unsimplified $(2-x)^{-1}$ expanded correctly, error in simplifying before their C is involved in the expression, allow A1FT when their C is introduced. The FT is on A, B, C . |
| | Multiply expansion of $\left(1 + \frac{3x^2}{2}\right)^{-1}$ (must reach $1 \pm \frac{3x^2}{2}$) by $Ax + B$, where $AB \neq 0$, up to the term in x^3 . Allow if used $Cx + D$ ($Ax + B$ miscopied). | M1 | Allow either ± 2 or $\pm 2^{-1}$ outside bracket or missing. Allow one error in actual multiplication to acquire the 4 terms [all terms needed]. Ignore errors in higher powers. |

| Question | Answer | Marks | Guidance |
|----------|--|-------------|---|
| (b) | Obtain final answer $4 + \frac{1}{4}x - \frac{13}{8}x^2 + \frac{29}{16}x^3$, or equivalent [If final answer has been multiplied throughout, e.g. by 16 then A0 at the end] | A1 5 | Maclaurin's Series: $f(0) = 4 \quad B1 \quad f'(0) = 1/4 \quad B1.$ $f''(0) = -13/4$ and $f'''(0) = 87/8 \quad B1.$ $4 + \frac{1}{4}x - \frac{13}{8}x^2 + \frac{29}{16}x^3$ or equivalent M1 A1. If $4 + \frac{1}{4}x - \frac{13}{8}x^2 + \frac{29}{16}x^3$ then M0 A0 unless their $f(0)$ actually is 1. |
| (c) | State answer $ x < \sqrt{\frac{2}{3}}$ or $-\sqrt{\frac{2}{3}} < x < \sqrt{\frac{2}{3}}$ clear conclusion required | B1 1 | Or exact equivalent. Strict inequality. |

23. 9709_m22_ms_32 Q: 1

| Question | Answer | Marks | Guidance |
|--|--|-----------|--|
| | State or imply non-modular inequality $(2x+3)^2 > 3^2(x+2)^2$, or corresponding quadratic equation, or pair of linear equations | B1 | |
| | Make a reasonable attempt at solving a 3-term quadratic, or solve two linear equations for x | M1 | Quadratic formula or $(5x+9)(x+3)$ |
| | Obtain critical values $x = -3$ and $x = -\frac{9}{5}$ | A1 | OE |
| | State final answer $-3 < x < -\frac{9}{5}$ or $x > -3$ and $x < -\frac{9}{5}$ | A1 | [Do not condone \leq for $<$ in the final answer.] No ISW |
| Alternative method for question 1 | | | |
| | Obtain critical value $x = -3$ from a graphical method, or by solving a linear equation or linear inequality | B1 | |
| | Obtain critical value $x = -\frac{9}{5}$ similarly | B2 | |
| | State final answer $-3 < x < -\frac{9}{5}$ or $x > -3$ and $x < -\frac{9}{5}$ | B1 | [Do not condone \leq for $<$ in the final answer.] No ISW |
| | | 4 | |

compiled by examinent.com

24. 9709_S22_ms_31 Q: 2

| Question | Answer | Marks | Guidance |
|----------|--|--------------|--|
| (a) | State a correct unsimplified version of the x^2 or the x^4 term of the expansion of $(2-x^2)^{-2}$ or $\left(1-\frac{1}{2}x^2\right)^{-2}$ | M1 | $\frac{1}{4}\left(1+2\frac{x^2}{2}+\frac{-2}{2}-3\left(\frac{x^2}{2}\right)^2\right)$ Symbolic binomial coefficients are not sufficient for the M1. |
| | State correct first term $\frac{1}{4}$ | B1 | Accept 2^{-2} . |
| | Obtain the next two terms $\frac{1}{4}x^2 + \frac{3}{16}x^4$ | A1 A1 | A1 for each one correct ISW. Full marks for $\frac{1}{4}(1+x^2 + \frac{3}{4}x^4)$ ISW. |
| | | | SC allow M1 A1 A1 for $\frac{1}{4}$ and $1+x^2 + \frac{3}{4}x^4$ SOI. SC allow M1 A1 for $1+x^2 + \frac{3}{4}x^4$ |
| | | 4 | |
| (b) | State answer $ x < \sqrt{2}$ | B1 | Or $-\sqrt{2} < x < \sqrt{2}$. |
| | | 1 | |

compiled by examinent.com

25. 9709_S22_ms_31 Q: 5

| Question | Answer | Marks | Guidance |
|---|---|-------|---|
| (a) | Substitute $x = 2$, equate to zero | M1 | Or divide by $x - 2$ and equate constant remainder to zero. |
| | Obtain a correct equation, e.g. $8a - 40 + 2b + 8 = 0$ | A1 | Seen or implied in subsequent work. |
| | Differentiate $p(x)$, substitute $x = 2$ and equate result to zero | M1 | Or divide by $x - 2$ and equate constant remainder to zero. |
| | Obtain $12a - 40 + b = 0$, or equivalent | A1 | SOI in subsequent work. |
| | Obtain $a = 3$ and $b = 4$ | A1 | |
| Alternative method for question 5(a) | | | |
| | State or imply $(x-2)^2$ is a factor | M1 | |
| | $p(x) = (x-2)^2(ax+2)$ | A1 | |
| | Obtain an equation in b | M1 | |
| | e.g. by comparing coefficients of x : $b = 4a - 8$ | A1 | |
| | Obtain $a = 3$ and $b = 4$ | A1 | |
| | | | SC If uses $x = -2$ in both equations allow M1 and allow A1 for $a = -3$, $b = -4$. |
| | | 5 | |

| Question | Answer | Marks | Guidance |
|---|---|-------|---|
| (b) | Attempt division by $(x - 2)$ | M1 | The M1 is earned if division reaches a partial quotient of $ax^2 + kx$, or if inspection has an unknown factor $ax^2 + ex + f$ and an equation in e and/or f . Where a has the value found in part 5(a). |
| | Obtain quadratic factor $3x^2 - 4x - 4$ | A1 | |
| | Obtain factorisation $(3x+2)(x-2)(x-2)$ | A1 | |
| Alternative method for question 5(b) | | | |
| | State or imply $(x-2)^2$ is a factor | B1 | |
| | Attempt division by $(x-2)^2$, reaching a quotient $ax + k$ or use inspection with unknown factor $cx + d$ reaching a value for c or for d | M1 | |
| | Obtain factorisation $(3x+2)(x-2)^2$ | A1 | Accept $3\left(x + \frac{2}{3}\right)(x-2)^2$. |
| | | 3 | |

To access the full eBook and explore more content, visit:

www.examinent.com